



WALKING POSTER PRESENTATION

Open Access

3D T1-weighted self-gated cardiac MRI for assessing myocardial infarction in mouse models

Xiaoyong Zhang^{1,2}, Hanwei Chen³, Zijun Wei², Fei Yan², Shi Su², Yanchun Zhu², Bensheng Qiu¹, Xin Liu^{2*}, Zhaoyang Fan⁴, Guoxi Xie²

From 19th Annual SCMR Scientific Sessions
 Los Angeles, CA, USA. 27-30 January 2016


Background

Mouse models with myocardial infarction (MI) have been intensively used to investigate the cardiac remodeling and functional change [1,2]. MRI with ECG triggering and respiratory gating (or breath-holding) is commonly used to assess MI due to its noninvasive nature. However, the manipulations of external ECG triggering and respiratory gating are cumbersome due to the small size of mouse. To address this issue, a 3D self-gating (SG) MR technique with stack-of-stars sampling trajectories was proposed for retrospectively cardiac and respiratory-gated MI imaging in mouse models [3].

Methods: Technical Design: MR data was acquired by a T1-weighted GRE sequence with stack-of-stars sampling trajectories and a partition-first golden-angle reordering (Figure 1a). The centers of k-space lines acquired at the same angle from individual partitions were aggregated and used as an SG time point. The periodic respiratory and cardiac motions were then detected through an iterative filtering process on the SG time series according to the cardiac rate of 300~500 per minute and respiration rate of 70~110 per minute. After resorting the imaging data into appropriate cardiac and respiratory phases, motion-artifact-free were finally reconstructed.

Experiments: The SG technique was preliminarily validated on 5 mice with MI induction and all MR scans were performed on a 3T scanner (Siemens Tim Trio, Germany) with a customized 4-channel mouse coil. Typical imaging parameters for the ungated GRE sequence included: flip angle = 18°, TR = 4.2 ms, TE = 2.4 ms, spatial resolution = 0.6 × 0.6 × 1.5 mm³, bandwidth = 620 Hz/Pixel, partition number = 12, and a

total number of 3200 projections were continuously collected, corresponding to a fixed scan time of 3 min. An amount of 0.5 ml gadolinium contrast agent with concentration of 0.5 mmol/ml (Consun Pharmaceutical Group Limited, GuangZhou, China) was injected to enhance the MI. The mice were sacrificed immediately after MRI for histological analysis and comparison to the MR results.



²Shenzhen Institutes of Advanced Technology, Shenzhen, China
 Full list of author information is available at the end of the article




Figure 2 (a) Representative histopathology pictures (top row) and reconstructed MR images (bottom row) from a mouse with MI induction. The regions of MI (arrows) matched nicely between the histology and the proposed SG technique. (b) The correlation of the ratio of the infarction size and myocardial size between histology and the proposed method.

Results

All MR scans were successfully conducted. The cardiac motion detected by the SG technique matched nicely to the ECG signals recorded by a small animal ventilator (Chengdu Taimeng Software Co.LTD, Chengdu, China) (Fig. 1b&1c). And the MI regions detected by the SG

technique were also matched well to the histopathology analysis results (Fig. 2).

Conclusions

A 3D SG technique was developed for assessing MI in mouse model. The preliminary in vivo study has

demonstrated that the technique can correctly detect the MI, which may outperform the conventional MR techniques with ECG-triggering and respiratory gating.

Authors' details

¹University of Science and Technology of China, Hefei, China. ²Shenzhen Institutes of Advanced Technology, Shenzhen, China. ³Guangzhou Panyu Central Hospital, Guangzhou, China. ⁴Cedars-Sinai Medical Center, Los Angeles, CA, USA.

Published: 27 January 2016

References

1. Lutgens E, et al: *Cardiovascular research* 1999, **41**(3):586-593.
2. Hofmann U, et al: *Circulation* 2012, **125**(13):1652-1663.
3. Liu J, et al: *MRM* 2010, **63**(5):1230-1237.

doi:10.1186/1532-429X-18-S1-Q49

Cite this article as: Zhang et al.: 3D T1-weighted self-gated cardiac MRI for assessing myocardial infarction in mouse models. *Journal of Cardiovascular Magnetic Resonance* 2016 **18**(Suppl 1):Q49.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

